



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

REVIEWS OF RECENT LITERATURE.

ANTHROPOLOGY.

Human Remains from Maya Territory.—A few broken calvaria, a few fragments of long bones, and a few sets of teeth are all the skeletal remains that we possess of the ancient inhabitants of the Maya territory. In the admirable report upon the ruins of Copan, by Maudsley, we read that "traces of bones were found," "a few small fragments of human bones were found," etc. The more recent explorations at the same ruin by the expeditions from the Peabody Museum have resulted in the discovery of "fragments" and "filed teeth." The few crania which, it might be inferred from the field notes, were well preserved and which are now in the Peabody Museum are in too fragmentary a condition to be available for study. In his account of the ruins of Xkichmook,¹ Mr. Edward Thompson describes the structure and architectural remains of an interesting group of ruins. We may readily believe that the "Palace," which he pictures in word, photograph, and plan, is "a most imposing structure," as its walls stand 64 feet high and are raised 80 feet above the surrounding level. Notwithstanding the probably dense population, as indicated by the extent of the ruins which formerly occupied the region, very few human remains were found. "Badly decayed human bones and teeth," "fragments of human bones much decayed," etc., is the record of the explorations among the burial chambers.

In a pleasantly written narrative² Mr. Gordon gives an account of the Honduras of to-day and of the vast quantities of potsherds and other artifacts that exist along the Uloa River. Whether or not they were manufactured by a race distinct from the Mayas, as suggested by Mr. Gordon and Professor Putnam in his editorial note, they indicate the presence of an extensive population in that valley in former times. However, the disappointing statement again appears, "The human remains . . . consist of crumbling fragments of bone . . .

¹ Thompson, Edward. Ruins of Xkichmook. Field Columbian Museum, *Anthropological Series*, vol. ii, No. 3.

² Gordon, George B. Researches in the Uloa Valley, Caverns of Copan, Honduras, *Memoirs of the Peabody Museum of American Archaeology and Ethnology*. Cambridge, Mass. Vol. i (1898), Nos. 4 and 5.

too minute to supply any information respecting the form of the burials," or, what is much more important, the character of the skeletons themselves, from which the racial affinities of the people might perhaps be determined. Scarcely second in interest to the long-sought "Rosetta stone of the Mayas" would be a moderately large series of Maya skeletons in a fair state of preservation.

FRANK RUSSELL.

GENERAL BIOLOGY.

Variation in Seedlings.—Professor Herbert L. Jones¹ found in Cambridge, Mass., a sycamore maple, *Acer pseudoplanatus*, which produces seedlings showing a large number of abnormalities. The cotyledons show a tendency to be doubled, and all grades of doubling were found from a mere notch at the distal end of one cotyledon to four complete seed-leaves. Where one cotyledon is completely doubled, the plumule usually consists of a whorl of three leaves, but where there are four complete cotyledons there are never more than two leaves—a very curious correlation.

R. P. B.

Chemical Analysis of the Plankton.—The chemical composition of the plankton of the Baltic Sea is discussed and compared with the analysis of certain agricultural products in a recent paper by Brandt.² The Copepoda, Peridinidæ, and diatoms are the predominant forms in the plankton of this body of water. Fifty cubic centimeters of plankton of mixed constitution weigh, when dried, from 1.08 to 1.74 grams; when diatoms predominate, the weight falls as low as .04 gram; and when Copepoda are abundant, it may rise as high as 2.1 grams. The results of the enumerations made of the organisms in the different catches subjected to analysis are utilized in the computation of their number in a gram of the dried plankton. It has thus been determined that it takes 675,000,000 diatoms (principally Chætoceros) or 42,000,000 to 65,000,000 Peridinidæ to weigh one gram; in the case of the marine Copepoda, from 300,000 to 500,000 are required; while the single analysis of a fresh-water plankton composed mainly of the larger individuals indicates that 50,000 to 124,000 are

¹ Jones, H. L. Unusual Forms of Maple Seedlings. Oberlin College, *Laboratory Bulletin* (1898), No. 9.

² Brandt, K. Beiträge zur Kenntniss der chemischen Zusammensetzung des Planktons. *Wissensch. Meeresuntersuchungen*, Neue Folge, Bd. iii (1898), Heft 2, pp. 45-90.